


Graphene enables ultrafast laser pulses for spectroscopy and biomedical applications

August 4, 2015



Schematic of an all-fibre laser capable of producing few-cycle pulses. Laser pulses of length 263 femtoseconds are produced from a seed oscillator (top) mode-locked using a graphene-based saturable absorber. These pulses are fed through a ... more

Graphene Flagship researchers have developed an optical fibre laser that emits pulses with durations equivalent to just a few wavelengths of the light used. This fastest ever device based on graphene will be ideal for use in ultrafast spectroscopy, and in surgical lasers that avoid heat damage to living tissue.

Amazing Skin

Grandma's Wrinkle Trick
Nobel Prize winning scientists have discovered a shocking natural skincare solution. If you're 35+, try this one weird trick & look years younger...

[Read More >>](#)

Ultrafast and ultrashort

Advanced photonics applications such as high-speed spectroscopy require ultrashort pulses in order to capture transient physical phenomena in the materials studied. In practice, that means laser pulses in the femtosecond (10-15s) range. An example of such an application is pump-probe spectroscopy of photochemical relaxation processes.

"When engineering light to travel in ultrashort pulses, it is important to understand its wave nature," says Daniel Popa, head of the photonics group at the Cambridge Graphene Centre, and leader of its graphene-based laser research project. "For light to propagate as does a mechanical wave on a stretched cord, the shortest possible pulse is

defined by a single wave oscillation."

Time resolution is limited by the length of the laser pulse used. The shorter the pulse, the higher the spectroscopic resolution, with the highest possible resolution defined by the cycle length of the particular light frequency employed. In the visible and near-infrared regimes, in which most ultrafast lasers operate, the ultimate pulse duration lies between 2 and 5 femtoseconds. Shorter pulses require shorter wavelengths.

Theoretical limits aside, pulses as short as two cycles can be generated from laser cavities using a technique known as passive mode-locking. With titanium-sapphire lasers, common in photonics laboratories the world over, pulses of 5 femtosecond length can be produced at a wavelength of 800 nanometres, corresponding to less than two cycles. These pulses are not tuneable, however. Tuneable few-cycle pulses can be achieved by exploiting nonlinear effects in optical parametric amplifiers, but the practical arrangements tend to be complex and expensive.

Fibre lasers are attractive platforms for ultrashort pulse generation, owing to their simple, compact and cost-effective designs, their efficient heat dissipation, and an alignment-free operation that does not require bulky optical setups. With fibre-based oscillators, ultrashort pulses can be generated by passive mode-locking, which requires a nonlinear component known as a saturable absorber. Graphene has the ideal physical properties to make such a saturable absorber.

Start Download Now

Convert From Doc to PDF, PDF to Doc Simply With The Free On-line App!

A graphene-based all-fibre laser for few-cycle light pulses

Graphene-based mode-locked lasers have been demonstrated before, but it is the use of this novel

Featured	Last comments	Popular
	Study shows bumblebees fly differently depending on the load they are carrying 9 hours ago	0
	Computer simulation shows climate mitigation schemes could result in increased water stress 8 hours ago	9
	Researchers suggest office thermostat settings biased against women 9 hours ago	4
	Thunderstrike 2: Proof-of-concept worm could infect Macs 12 hours ago	1
	For faster battery charging, try a quantum battery? Aug 03, 2015	6

[more »](#)

Phys.org

Segui

+114.056

Phys.org on facebook

Like Sandro Mignuzzi, Ahmed Kiani and 1,068,195 others like this.

Relevant PhysicsForums posts

- Recognize this irreversible equation for kinetic modeling? 1 hour ago
- Thermally insulating tape at high temperature (350C) 12 hours ago
- Difference between Creep and Strain Rate Sensitivity 18 hours ago
- CO2 laser recirculation compressors and coolers 21 hours ago
- Carbon nanotube exoskeleton Aug 03, 2015
- Thermochromic Materials for IR absorption Aug 03, 2015
- More from Materials and Chemical Engineering

and compression of the pulses.

The key is to extract the pulse from such a cavity when its duration is at a minimum, and peak power thus at a maximum. Owing to the high peak power of the extracted pulse, new frequency components can be generated through nonlinear optical effects within an external length of fibre, and these are critical when it comes to further decreasing the pulse length. This is based on the mathematical relationship in waves between frequency and time domains known as a Fourier transform. To realise this transformation in physical form, the researchers engineered a dispersive delay line that folds the newly created frequency components into a single pulse.

The Graphene Flagship researchers' setup was based only on standard telecommunications equipment, with a saturable absorber based on a composite of graphene and polyvinyl alcohol (PVA) fabricated by low-cost solution processing, with the graphene flakes exfoliated from bulk graphite by ultrasonic agitation of the solution. Evaporation leaves behind a 50 micron-thick graphene-PVA composite, which is then sandwiched between fibre connectors.

With this setup, Purdie and his colleagues were able to generate 29 femtosecond pulses, which corresponds to fewer than six cycles at a wavelength of 1.5 microns.

Compensating for higher-order nonlinear and dispersive effects should lead to a shorter pulse length, and the use of a higher power diode, or a double-pumped configuration, could result in higher bandwidth pulses as well as increased output power. Finally, the addition of photonic crystal fibres could in principle allow for the generation of similarly short laser pulses at other wavelengths.

"What is truly remarkable about this project is the ease of combining graphene with off-the-shelf optical fibres in a highly compact format," says Popa. "In this way, we can generate light pulses that last for only a few cycles, or a few millionths of a billionth of a second."

Explore further: The future of ultrashort laser pulses

More information: "Few-cycle pulses from a graphene mode-locked all-fiber laser." *Appl. Phys. Lett.* 106, 253101 (2015); [dx.doi.org/10.1063/1.4922397](https://doi.org/10.1063/1.4922397)

Journal reference: [Applied Physics Letters](#)

0 shares

Provided by: Graphene Flagship

[feedback to editors](#)



15 Of The Rarest (And Most Mind Blowing) Photographs In History
LOLWOT



Man Who Explored Empty Silo Wishes He Hadn't
HyperTomb



Abandoned Luxury Cars from Dubai
Lists and News



10 of The Creepiest Photographs Ever Taken
Time To Break


Sponsored Links by Taboola

Related Stories



The future of ultrashort laser pulses July 24, 2014

Rapid advances in techniques for the creation of ultra-short laser pulses promise to boost our knowledge of electron motions to an unprecedented level.



New method to generate arbitrary optical pulses January 21, 2015

Scientists from the University of Southampton have developed a new technique to generate more powerful, more energy efficient and low-cost pulsed lasers.



New technology makes creating ultrashort infrared laser pulses easy and cheap January 27, 2015

In a marathon, everyone starts at roughly the same place at roughly the same time. But

Recommended for you



Fast times and hot spots in plasmonic nanostructures

August 4, 2015
The ability to control the time-resolved optical responses of hybrid plasmonic nanostructures was demonstrated by a team led by scientists in the Nanophotonics Group at the Center for Nanoscale Materials including collaborators ...



New clot-busting treatment targets number one killer August 4, 2015

Australian researchers funded by the National Heart Foundation are a step closer to a safer and more effective way to treat heart attack and stroke via nanotechnology.



Study explores nanoscale structure of thin films August 4, 2015

The world's newest and brightest synchrotron light source—the National Synchrotron Light Source II (NSLS-II) at the U.S. Department of Energy's Brookhaven National

User comments

Please [sign in](#) to add a comment. Registration is free, and takes less than a minute. [Read more](#)

[Click here to reset your password.](#)
Sign in to get notified via email when new comments are made.

